

PATCH TRANSDERMIQUE

FICHE N° 9530

PRÉSERVER
SAUVEGARDER
VALORISER

Période de fabrication : -

Fabricant : fabricant non renseigné

Domaines : Santé

Sous-domaines :

Organisme : Université d' Auvergne, Faculté de médecine

Ville : Clermont-Ferrand

Modèle :

Matériaux :

Description

Patch de forme ronde, autocollant, conservé dans un emballage sous forme de sachet.

Utilisation

Le patch se colle sur le bras. Il est utilisé comme remède pour traiter les angines.



1 Pad 16 cm² • No. 581
NDC 0025-2068

NitrodiscTM

32 mg
(nitroglycerin)

Caution: Federal law prohibits dispensing without prescription.

Usual Adult Dose: Apply one pad to skin daily. See accompanying complete prescribing information.

Store at controlled room temperature of 59°-86°F (15°-30°C). Do not refrigerate. Avoid extremes of temperature and/or humidity.

SEARLE

DIRECTIONS FOR USE



1. Tear pouch open starting at corner as shown. Remove Nitrodisc.TM



2. Peel protective foil covering from Nitrodisc.



3. • Apply Nitrodisc to site free of hair on chest or on inner arm.
• Change application site each time to avoid skin irritation.
• If the Nitrodisc falls off or loosens, a new Nitrodisc should be applied.
• Apply a new Nitrodisc every 24 hours or after bathing.

Searle Pharmaceuticals Inc.
Chicago IL 60680

103744-1

AO5250-1

INSTRUCTIONS FOR THE PATIENT

NitrodiscTM
(nitroglycerin)

Once-a-Day Therapy for Angina Pectoris

Nitrodisc — How It Works

Nitrodisc is a timed-release nitroglycerin product. Nitroglycerin or related medications are available in several forms. Sublingual tablets, which are to be placed under the tongue, act quickly, but they have an effect for only a short period of time. Long-acting

©1981, Searle Pharmaceuticals Inc.

1 Pad 8 cm² • No. 580
NDC 0025-2058

NitrodiscTM
16 mg
(nitroglycerin)

Caution: Federal law prohibits dispensing without prescription.

Usual Adult Dose: Apply one pad to skin daily. See accompanying complete prescribing information.

Store at controlled room temperature of 59°-86°F (15°-30°C). Do not refrigerate. Avoid extremes of temperature and/or humidity.

SPARLE

headaches, your doctor may adjust your dosage and/or recommend that you also take headache medication.



- Although uncommon, faintness, flushing, weakness, or dizziness may occur, especially when suddenly arising from a lying, sitting, or bending position. Nausea, vomiting, and skin rash are other uncommon side effects. If any of these persist, you should notify your physician.

- If your angina attacks change for the worse—if they become more severe, occur more frequently, last longer, or change in any way—advise your physician promptly.

Revised: May 12, 1981

SEARLE

Searle Pharmaceuticals Inc.
Box 5110, Chicago IL 60680

Printed in USA

Pour nous citer :

Base de la Mission nationale de sauvegarde et de valorisation du patrimoine scientifique et technique contemporain, PATSTEC, Patch transdermique (fabricant non renseigné), <https://www.patstec.fr/ressources/objets/detail?id=14141>, consulté le 2026-07-27